

Nuclear Stress Test Instructions for Procedure at Eastern Cardiology, P.A.

11	eadmil	1:	1	Day		2 Day							
DA	ATE:								_ TIM	E:			
You	u musi					least 2- you wi						<u>l your</u>	<u>-</u>
1. <b>D</b>	O NO	T	lrin	k an	y co	ffee, t	ea,	soda	s, die	t drin	ks o	eat	any
					•	cludin	,		,				•
VO	ur pr	oce	edur	e.									
•	_				omfo	rtable sh	oes a	and lo	ose-fittii	ng cloth	ning. F	emale 1	oatients
						fit, no f				_	_	-	-
	lognes			•				•	•			•	
3. YC	M LIC	<b>37</b> T	A TZT	ATT	O D 37	OTTD 3.1	~~.		CODITA	70 1 10	DICAT	77.07.70	
٦. ١	JU MIA	YIA	AKE	ALL (	OF Y	OUR NO	ORM	IAL N	IORNIN	IG ME	DICA.	TONS	except
	e follow			ALL (	OF Y	OUR NO	ORN	IAL N	10RNIN	NG MEI	DICA .	TONS	except
	e follow	ing:				OUR NO							except
the	e follow ke for 2	ing: 4 ho		BL			SURI			ΓED BI	ELOW		
the Do not tal	e follow ke for 2	ing: 4 ho Ate	ours	BL		PRESS	SURI rolol		DS LIS	ΓED BI	ELOW Ca	7	ol
Do not tal	e follow ke for 2 ol	ing: 4 ho Ate Ind	ours enolol leral	BL	OOD	PRESS Bisopr Imdur	SURI colol		DS LIST	ΓΕD BI ic bide	ELOW Ca La	, arvedil	ol l
Do not tal Acebutolo Coreg	e follow ke for 2 ol	ing: 4 ho Ate Ind Me	ours enolol leral	BL olol/To	OOD	PRESS Bisopr Imdur	SURI colol		DS LIST Bystoli Isosori	FED BI ic bide blol	ELOW Ca La Pr	rvedil abetalo	ol l
Do not tal Acebutolo Coreg Lopressor	e follow ke for 2 bl	ing: 4 ho Ate Ind Met	ours enolol leral topro normi	BL olol/To	OOD	PRESS Bisopr Imdur Nadole	SURI colol		DS LIST Bystoli Isosori Nebivo	FED BI ic bide blol	ELOW Ca La Pr	rvedil abetalo opran	ol l
Do not tal Acebutolo Coreg Lopressor Sectral Do not tal	e follow ke for 2 ol c ke for 4	ing: 4 ho Ate Ind Met	ours enolol leral topro normi	BL olol/To	OOD	PRESS Bisopr Imdur Nadole	SURI colol colol ate	E MEI	DS LIST Bystol Isosor Nebivo Visker	FED BI ic bide blol	ELOW Ca La Pr	rvedil abetalo opran	ol l
Do not tal Acebutolo Coreg Lopressor Sectral	e follow ke for 2 ol c ke for 4	Ate Ate Ind Met Ten 8 ho	ours enolol leral topro normi	BL olol/To	OOD	PRESS Bisopr Imdur Nadole	SURI colol colol ate		DS LIST Bystol Isosor Nebivo Visker	FED BI ic bide blol	ELOW Ca La Pr	rvedil abetalo opran	ol l
Do not tal Acebutolo Coreg Lopressor Sectral Do not tal Aggrenox	e follow ke for 2 ol c ke for 4	Ate Ind Mer Ten 8 ho	ours enolol leral topro normi	BL	oprol	PRESS Bisopr Imdur Nadole	SURI colol colol ate	E MEI	DS LIST Bystoli Isosori Nebivo Visken	FED BI ic bide blol	ELOW Ca La Pr Zo	rvedil abetalo opran	ol ol olol

- you may have a light breakfast (toast, cereal, oatmeal, etc) before 8am.
- 5. Please bring 2 snacks (pack of crackers, a sandwich, fruit, juice beverage, etc.). You will eat 2 times during the test. DO NOT EAT UNTIL THE TECHNICIAN ADVISES YOU.
- 6. Allow 4 to 5 hours in our office for the test to be completed including testing and resting time.
- 7. If you are having a 2-day nuclear please allow 2 hours in our office each day.
- 8. Do not smoke the morning of the test

## PLEASE FOLLOW ALL INSTRUCTIONS AS DIRECTED